

Pseudoholomorphic discs near an elliptic point

Alexandre Sukhov* and Alexander Tumanov**

*Université des Sciences et Technologies de Lille, Laboratoire Paul Painlevé, U.F.R. de Mathématique, 59655 Villeneuve d'Ascq, Cedex, France , e-mail: sukhov@math.univ-lille1.fr

** Department of Mathematics, University of Illinois, 1409 West Green Street, Urbana, IL 61801, USA, e-mail: tumanov@math.uiuc.edu

Abstract. We prove the existence and study the geometry of Bishop discs near an elliptic point of a real n -dimensional submanifold of an almost complex n -dimensional manifold.

MSC: 32H02, 53C15.

Key words: almost complex structure, generic manifold, elliptic point, Bishop disc.

1 Introduction

E.Bishop [2] proved that holomorphic discs with boundaries attached near an elliptic point p of a real n -dimensional submanifold E of $\mathbb{C}^n$ form a family smoothly depending on $(n-1)$ real parameters and that the boundaries of these discs fill a neighborhood of p in E . A substantial progress in the study of the geometry of these discs (called Bishop discs) has been done by E.Bedford - B.Gaveau [1], C.Kenig - S.Webster [6, 7] and other authors. On the other hand, M.Gromov in [4] pointed out that Bishop discs still exist when E is a submanifold of an almost complex manifold; he used this result in his approach to the symplectic and contact geometry. In the case of almost complex dimension 2, a precise version of this result was established and substantially used by H.Hofer [5], Y.Eliashberg [3], R.Ye [12] in various forms. The goal of the present paper is to give a complete proof of the existence of Bishop discs with boundaries attached near an elliptic point p of a real n -dimensional submanifold of an almost complex manifold of complex dimension n . We prove that their boundaries fill a neighborhood of p on E similarly to the case of the standard complex structure. Our approach arises from the standard complex analysis techniques, in particular, we use the methods of the works [8] and [10].

2 Preliminaries

2.1 Almost complex manifolds.

Let (M, J) be an almost complex manifold with operator of complex structure J . Let $\mathbb{D}$ be the unit disc in $\mathbb{C}$ and J_{st} the standard (operator of) complex structure on $\mathbb{C}^n$ for arbitrary n . Let f be a smooth map from $\mathbb{D}$ into M . We say that f is J -holomorphic if $df \circ J_{st} = J \circ df$. We call such a map f a J -holomorphic disc and denote by $\mathcal{O}_J(\mathbb{D}, M)$ the set of J -holomorphic discs in M . We denote by $\mathcal{O}(\mathbb{D})$ the space of usual holomorphic functions on $\mathbb{D}$.

The following lemma shows that an almost complex manifold (M, J) can be locally viewed as the unit ball $\mathbb{B}$ in $\mathbb{C}^n$ equipped with a small almost complex deformation of J_{st} . We shall repeatedly use this observation in what follows.

Lemma 2.1 *Let (M, J) be an almost complex manifold. Then for each $p \in M$, each $\delta_0 > 0$, and each $k \geq 0$ there exist a neighborhood U of p and a smooth coordinate chart $z : U \rightarrow \mathbb{B}$ such that $z(p) = 0$, $dz(p) \circ J(p) \circ dz^{-1}(0) = J_{st}$, and the direct image $z_*(J) := dz \circ J \circ dz^{-1}$ satisfies the inequality $\|z_*(J) - J_{st}\|_{C^k(\mathbb{B})} \leq \delta_0$.*

Proof : There exists a diffeomorphism z from a neighborhood U' of $p \in M$ onto $\mathbb{B}$ such that $z(p) = 0$ and $dz(p) \circ J(p) \circ dz^{-1}(0) = J_{st}$. For $\delta > 0$ consider the isotropic dilation $d_\delta : t \mapsto \delta^{-1}t$ in $\mathbb{C}^n$ and the composite $z_\delta = d_\delta \circ z$. Then $\lim_{\delta \rightarrow 0} \|(z_\delta)_*(J) - J_{st}\|_{C^k(\mathbb{B})} = 0$. Setting $U = z_\delta^{-1}(\mathbb{B})$ for sufficiently small positive δ we obtain the required result.

The operators ∂_J and $\bar{\partial}_J$.

Let (M, J) be an almost complex manifold. We denote by TM the real tangent bundle of M and by $T_{\mathbb{C}}M$ its complexification. Recall that $T_{\mathbb{C}}M = T^{(1,0)}M \oplus T^{(0,1)}M$ where $T^{(1,0)}M := \{X \in T_{\mathbb{C}}M : JX = iX\} = \{\zeta - iJ\zeta, \zeta \in TM\}$, and $T^{(0,1)}M := \{X \in T_{\mathbb{C}}M : JX = -iX\} = \{\zeta + iJ\zeta, \zeta \in TM\}$. Let T^*M be the cotangent bundle of M . Identifying $\mathbb{C} \otimes T^*M$ with $T_{\mathbb{C}}^*M := \text{Hom}(T_{\mathbb{C}}M, \mathbb{C})$ we define the set of complex forms of type $(1, 0)$ on M as $T_{(1,0)}M = \{w \in T_{\mathbb{C}}^*M : w(X) = 0, \forall X \in T^{(0,1)}M\}$ and we denote the set of complex forms of type $(0, 1)$ on M by $T_{(0,1)}M = \{w \in T_{\mathbb{C}}^*M : w(X) = 0, \forall X \in T^{(1,0)}M\}$. Then $T_{\mathbb{C}}^*M = T_{(1,0)}M \oplus T_{(0,1)}M$. This allows us to define the operators ∂_J and $\bar{\partial}_J$ on the space of smooth functions on M : for a smooth complex function u on M we set $\partial_J u = du_{(1,0)} \in T_{(1,0)}M$ and $\bar{\partial}_J u = du_{(0,1)} \in T_{(0,1)}M$. As usual, differential forms of any bidegree (p, q) on (M, J) are defined by exterior multiplication.

2.2 Real submanifolds of an almost complex manifold

Let E be a real submanifold of codimension k in an almost complex manifold (M, J) of complex dimension n . Locally E is defined as the zero-set of a smooth $\mathbb{R}^k$ -valued function $r = (r_1, \dots, r_k)$ with $dr_1 \wedge \dots \wedge dr_k \neq 0$. As usual, we denote by $H_p E$ the maximal complex subspace of the tangent space $T_p E$ that is $H_p E = T_p E \cap J(p)T_p E$ and call it the holomorphic or the complex tangent space of E at p . Recall that E is called generic at p if the complex dimension of $H_p E$ is equal to $n - k$. A real submanifold E is called totally real at p if $H_p E = \{0\}$. In particular, any generic submanifold of real codimension n is totally real.

Let E be a real n -dimensional submanifold in an almost complex manifold (M, J) . Our considerations are local so we simply indentify M with $\mathbb{R}^{2n}$ (or $\mathbb{C}^n$) and view J as a smooth matrix-valued function. In this paper we study the geometry of E near a point $p \in E$ admitting a tangent vector $X \in T_p E$ such that $J(p)X \in T_p E$, that is a complex tangent direction that is p is a singular point for the Cauchy-Riemann structure of E . We consider generic singularities only that is the case where there exists precisely one tangent complex line in $T_p E$. Our approach is based on a suitable choice of coordinates near p . First we can assume that $p = 0$ and $J(0) = J_{st}$. Let vector $e_1 = (1, 0, \dots, 0)$ be complex tangent to E at the origin. It follows by the classical Nijenhuis-Woolf theorem [9] that there exists a J -holomorphic disc $f : \mathbb{D} \rightarrow \mathbb{R}^{2n}$ such that $f(0) = 0$ and $df(0)(\frac{\partial}{\partial(\text{Re}\zeta)}) = e_1$. After a local diffeomorphism with the identical linear part at the origin straightening f , we obtain that the map $\zeta \mapsto (\zeta, 0, \dots, 0)$ is J -holomorphic on $\mathbb{D}$. Denote the coordinates in $\mathbb{C}^n$ by $Z = (w, z)$ with $z = (z_2, \dots, z_n)$. Then we may assume (possibly, after an additional change of coordinates preserving

the J -holomorphicity of the coordinates line w and a suitable choice of defining functions similarly to [2]) that E is defined near the origin by the equations

$$\begin{aligned}\rho(Z) &= z_2 - P(w) + R(Z) = 0, \\ r_3(Z) &= \operatorname{Re} z_3 + h_3(Z) = 0, \\ &\dots \\ r_n(Z) &= \operatorname{Re} z_n + h_n(Z) = 0\end{aligned}$$

where $R = O(|Z|^3)$ and $h_j(Z) = O(|Z|^2)$. Here

$$P(w) = w\bar{w} + \gamma(w^2 + \bar{w}^2)$$

The point p is called *elliptic* if in such a system of coordinates $\gamma \in [0, 1/2[$. This notion is invariant. In what follows we denote by $r := (\operatorname{Re} \rho, \operatorname{Im} \rho, r_3, \dots, r_n)$ the $\mathbb{R}^n$ -valued local defining function of E .

3 Bishop discs

Let (M, J) be a smooth almost complex manifold of real dimension $2n$ and E a real submanifold of M of real codimension m . A J -holomorphic disc $f : \mathbb{D} \rightarrow M$ continuous on $\overline{\mathbb{D}}$ is called a *Bishop disc* if $f(b\mathbb{D}) \subset E$ (where $b\mathbb{D}$ denotes the boundary of $\mathbb{D}$). Our aim is to prove the existence and to describe certain classes of Bishop discs attached to E . In this section we derive Riemann-Hilbert type boundary problems describing pseudoholomorphic Bishop discs near generic and elliptic points of a real submanifold in an almost complex manifold.

Since our considerations are local, we identify an almost complex manifold with $\mathbb{C}^n$ and denote by $Z = (Z_1, \dots, Z_n)$ the standard complex coordinates. An almost complex structure J may be viewed as a smooth real $(2n \times 2n)$ -matrix valued function $J : Z \mapsto J(Z)$ on a neighborhood of the origin in $\mathbb{C}^n$. We also may assume that $J(0) = J_{st}$. Then a smooth map

$$Z : \zeta \mapsto Z(\zeta)$$

$$Z : \mathbb{D} \rightarrow U$$

where U is a small enough neighborhood of the origin in $\mathbb{C}^n$ is J -holomorphic if and only if it satisfies the following PDE system

$$Z_{\bar{\zeta}} - A(Z)\overline{Z_{\zeta}} = 0$$

where $A(Z)$ is the complex $n \times n$ matrix of the operator whose composite with complex conjugation is equal to the endomorphism $(J_{st} + J_{\delta}(Z))^{-1}(J_{st} - J_{\delta}(Z))$ (which is an anti-linear operator with respect to the standard structure J_{st}). By lemma 2.1 we can assume that the norm of A is small enough.

3.1 Bishop discs and Bishop's equation near a generic point

Denote by Kf the Cauchy integral of a function f :

$$K(f)(\zeta) = \frac{1}{2\pi i} \int_{b\mathbb{D}} \frac{f(\tau)d\tau}{\tau - \zeta}$$

for $\zeta \in \mathbb{D}$ and by K_0f its principal value at boundary point:

$$K_0(f)(\zeta) = (v.p.) \frac{1}{2\pi i} \int_{b\mathbb{D}} \frac{f(\tau)d\tau}{\tau - \zeta}, \zeta \in \partial\mathbb{D}$$

We will use also the following notation for the Cauchy-Green transform:

$$T(f)(\zeta) = \frac{1}{2\pi i} \int \int_{\mathbb{D}} \frac{f(\tau)d\tau \wedge d\bar{\tau}}{\tau - \zeta}$$

Then for any function f of class C^1 on $\overline{\mathbb{D}}$ (the restriction C^1 can be replaced by some Sobolev's class, see [11]) we have the generalized Cauchy formula:

$$f = Kf + Tf_{\bar{\zeta}} \quad (1)$$

Denote also by $K_+f(\zeta)$ and $K_-f(\zeta)$ the limiting values of Kf at $\zeta \in b\mathbb{D}$ from inside and outside respectively.

Then we have the classical Plemelj-Sokhotski formulae (at every point of $b\mathbb{D}$):

$$K_{\pm}f = K_0f \pm (f/2), \quad (2)$$

and

$$K_+f - K_-f = f \quad (3)$$

In particular, (1) and (3) imply that

$$-Tf(\zeta) = K_-f(\zeta), \zeta \in b\mathbb{D} \quad (4)$$

Recall also for any real function $u \in L^2(b\mathbb{D})$ there exists a unique $v \in L^2(\overline{\mathbb{D}})$ (the Hilbert transform of u) such that $u + iv$ is a boundary value of a holomorphic function h from the Hardy space $H^2(b\mathbb{D})$. The Hilbert transform may be written as a singular integral operator:

$$Hf(\zeta) = \frac{1}{2\pi i} (v.p.) \int_0^{2\pi} \frac{e^{i\theta} + \zeta}{e^{i\theta} - \zeta} f(e^{i\theta}) d\theta$$

The classical fact is that for a non-integer $k > 0$, H is a continuous linear operator on $C^k(b\mathbb{D})$. Denote also by P_0f the value of the Poisson integral of f at the origin:

$$P_0f = \frac{1}{2\pi} \int_0^{2\pi} f(e^{i\theta}) d\theta$$

Integrating the identity $2\tau(\tau - \zeta)^{-1} = (\tau + \zeta)(\tau - \zeta)^{-1} + 1$ we obtain that

$$K_0 f = (i/2)Hf + (1/2)P_0 f$$

on the boundary of $\mathbb{ID}$. Now (2) gives

$$f(\zeta) = iHf(\zeta) + P_0 f - 2K_- f(\zeta), \zeta \in b\mathbb{ID} \quad (5)$$

Suppose that f is of class $C^\alpha(\overline{\mathbb{ID}})$ with $\alpha > 0$ non-intergral. Let also $f^* = u^* + iv^*$ be a function on $b\mathbb{ID}$ of class C^α .

Lemma 3.1 *Assume that f satisfies the following system of singular integral equations:*

$$f = Kf^* + Tf_{\bar{\zeta}}, \quad \zeta \in \mathbb{ID} \quad (6)$$

$$v^* = Hu^* + P_0 v^* + 2\text{Im}Tf_{\bar{\zeta}}, \quad \zeta \in b\mathbb{ID} \quad (7)$$

Then $f^* = f|_{b\mathbb{ID}}$.

Proof : (6) implies that $K_+ f^* = K_+ f$. In view of (4) we obtain $K_+ f = Kf^*$. Furthermore, (7) implies $-2\text{Im}Tf_{\bar{\zeta}} = \text{Im}(-v^* + Hu^* + P_0 v^*) = \text{Im}(iHf^* + P_0 f^* - f^*) = 2\text{Im}K_- f^*$. Therefore, $K_f = K_- f^*$ (recall that the value of the Cauchy type integral at the infinity is equal to 0). Thus, $K_+ f = K_+ f^*$ and $K_- f = K_- f^*$ which implies the statement of lemma.

Now we may write the Bishop equation. We assume that a real submanifold E is given by the following equations near the origin:

$$y = h(x, w)$$

with

$$h(0, 0) = 0, dh(0, 0) = 0,$$

where we use the notation $Z = (z, w) \in \mathbb{C}^{n-m} \times \mathbb{C}^m$, $z = x + iy$, $w = u + iv$. We also write a map Z in the form $Z(\zeta) = (z(\zeta), w(\zeta))$. We will represent the matrix A in the form

$$A = \begin{pmatrix} P & R \\ L & N \end{pmatrix}.$$

with $(m \times m)$ -bloc P .

Then a smooth disc Z is J -holomorphic and its boundary is attached to E if and only if it satisfies the following system of singular integral equations:

$$(RH_0) : \left\{ \begin{array}{l} z = Kz^* - T(P(Z)\bar{z}_{\bar{\zeta}} + R(Z)\bar{w}_{\bar{\zeta}}), \quad \zeta \in \mathbb{ID} \\ y^* = Hh(y^*, w^*) + y_0 - 2\text{Im}T(P(Z)\bar{z}_{\bar{\zeta}} + R(Z)\bar{w}_{\bar{\zeta}}), \quad \zeta \in b\mathbb{ID} \\ w = Kw^* - T(L(Z)\bar{z}_{\bar{\zeta}} + N(Z)\bar{w}_{\bar{\zeta}}), \quad \zeta \in \mathbb{ID} \\ v^* = Hu^* + v_0 - 2\text{Im}T(L(Z)\bar{z}_{\bar{\zeta}} + N(Z)\bar{w}_{\bar{\zeta}}), \quad \zeta \in b\mathbb{ID} \end{array} \right\}$$

where $Z^* = Z|_{b\mathbb{D}}$. Thus, this non-linear boundary Riemann-Hilbert type problem may be viewed as an almost complex analog of the standard Bishop equation. Applying the implicit function theorem we easily obtain that given non-integral $\alpha > 0$, real constants y_0, v_0 and a real function $u^* \in C^\alpha(b\mathbb{D})$ there exists a unique solution of the system (RH_0) .

We point out that the existence of Bishop discs attached to a generic submanifold of an almost complex manifold was proved in [8] by a different method. The present method is substantially more explicit and allows to write Bishop's equation as a singular integral equation similarly to the integrable case. This leads to a more convenient and explicit parametrization of the Banach space of Bishop's discs and can be useful for other applications. In particular, it follows immediately from the equation (RH_0) that Bishop's discs depend smoothly on deformations of the almost complex structure J .

3.2 Bishop discs and Bishop's equation near an elliptic point

Consider a real n -dimensional submanifold E in an almost complex n -dimensional manifold (M, J) and let p be an elliptic point of E . We assume without loss of generality that E is a submanifold in $\mathbb{C}^n$ and the origin is an elliptic point of E ; moreover, $J(0) = J_{st}$. Furthermore, choosing local coordinates similarly to the previous section, we assume that the map $\zeta \mapsto (\zeta, 0, \dots, 0)$ is J -holomorphic on $\mathbb{D}$. Denote as above the coordinates in $\mathbb{C}^n$ by $Z = (w, z)$ with $z = (z_2, \dots, z_n)$. Then we may assume that E is defined near the origin by the equations

$$\begin{aligned}\rho(Z) &= z_2 - P(w) + R(Z) = 0, \\ r_3(Z) &= \operatorname{Re} z_3 + h_3(Z) = 0, \\ &\dots \\ r_n(Z) &= \operatorname{Re} z_n + h_n(Z) = 0\end{aligned}$$

where $R = O(|Z|^3)$ and $h_j(Z) = O(|Z|^2)$. Here

$$P(w) = w\overline{w} + \gamma(w^2 + \overline{w}^2)$$

and $\gamma \in [0, 1/2[$. As above, we denote by $r := (\operatorname{Re} \rho, \operatorname{Im} \rho, r_3, \dots, r_n)$ the $\mathbb{R}^n$ -valued local defining function of E .

Then a smooth map f defined on $\mathbb{D}$ and continuous on $\overline{\mathbb{D}}$ is a Bishop disc if and only if it satisfies the following non-linear boundary problem of the Riemann-Hilbert type for the quasi-linear operator $\overline{\partial}_J$:

$$(RH) : \begin{cases} \overline{\partial}_J f(\zeta) = 0, \zeta \in \mathbb{D} \\ r(f)(\zeta) = 0, \zeta \in b\mathbb{D} \end{cases}$$

Consider the non-isotropic dilations $\Lambda_\delta : Z = (w, z) \mapsto Z' = (\delta^{-1/2}w, \delta^{-1}z)$. In the new Z -variables (we drop the primes) the image $E_\delta = \Lambda_\delta(E)$ is defined by the equation $r_\delta(Z) := \delta^{-1}r((\Lambda_\delta)^{-1}Z) = 0$. Then the vector-function r_δ tends to the function

$$r^0(Z) = (\operatorname{Re} z_2 - \operatorname{Re} P(w), \operatorname{Im} z_2 - \operatorname{Im} P(w), \operatorname{Re} z_3 + Q_3(w), \dots, \operatorname{Re} z_n + Q_n(w))$$

as $\delta \rightarrow 0$. Here Q_j are some real homogeneous of degree 2 polynomials in w . Hence the manifolds E_δ approach the model quadric manifold $E_0 = \{r^0(Z) = 0\}$ which explicitly is given by the equations

$$\begin{aligned}\rho^0(Z) &= z_2 - P(w) = 0, \\ r_3^0(Z) &= \operatorname{Re} z_3 - Q_3(w) = 0, \\ &\dots \\ r_n^0(Z) &= \operatorname{Re} z_n - Q_n(w) = 0\end{aligned}$$

Consider the transported structures $J_\delta := (\Lambda_\delta)_*(J) = d\Lambda_\delta \circ J \circ (d\Lambda_\delta)^{-1}$. The following statement is similar to [8].

Lemma 3.2 *For any positive k and any compact subset $K \subset \mathbb{C}^n$ we have $\|J_\delta - J_{st}\|_{C^k(K)} \rightarrow 0$ as $\delta \rightarrow 0$.*

Proof : Consider the Taylor expansion of $J(Z)$ near the origin: $J(Z) = J_{st} + L(Z) + R(Z)$ where $L(Z)$ is the linear part of the expansion and $R(Z) = O(|Z|^2)$. Clearly, $\Lambda_\delta \circ R(\Lambda_\delta^{-1}(Z)) \circ \Lambda_\delta^{-1}$ converges to 0 as $\delta \rightarrow 0$. Denote by $L_{kj}^\delta(Z)$ (respectively, by $L_{kj}(Z)$) an entry of the real matrix $\Lambda_\delta \circ L(\Lambda_\delta^{-1}(Z)) \circ \Lambda_\delta^{-1}$ (respectively, of $L(Z)$). For $k = 3, \dots, 2n$, $j = 1, 2$ we have $J_{kj}^\delta(w, z) = \delta^{-1/2} L_{kj}(\delta^{1/2} w, \delta z) \rightarrow L_{kj}(w, 0)$ as $\delta \rightarrow 0$. However, in the coordinate system fixed above the map $\zeta \rightarrow (\zeta, 0, \dots, 0)$ is J -holomorphic, so $L_{kj}(w, 0) = 0$ for $k, j = 1, 2$. This implies that L_{kj}^δ approaches 0 for all k, j (for other values of k, j this is obvious). This gives us the result of the lemma.

Consider a J_δ -holomorphic disc $Z : \mathbb{D} \rightarrow (\mathbb{B}^n, J_\delta)$. The J_δ -holomorphy condition $J_\delta(Z) \circ dZ = dZ \circ J_{st}$ can be written in the following form:

$$Z_{\bar{\zeta}} - A_{J,\delta}(Z) \overline{Z_{\bar{\zeta}}} = 0 \quad (8)$$

where as above $A_{J,\delta}(Z)$ is the complex $n \times n$ matrix of the operator whose composite with complex conjugation is equal to the endomorphism $(J_{st} + J_\delta(Z))^{-1}(J_{st} - J_\delta(Z))$ (which is an anti-linear operator with respect to the standard structure J_{st}). Hence the entries of the matrix $A_{J,\delta}(Z)$ are smooth functions of δ, Z vanishing identically in Z for $\delta = 0$.

Using the Cauchy-Green transform T_{CG} defined above we may write $\bar{\partial}_J$ -equation (8) as follows:

$$[Z - T_{CG} A_{J,\delta}(Z) \overline{Z_{\bar{\zeta}}}]_{\bar{\zeta}} = 0$$

This is well known [11] that the Cauchy-Green transform is a continuous linear operator from $C^k(\overline{\mathbb{D}})$ into $C^{k+1}(\overline{\mathbb{D}})$ (recall that k is nonintegral). Hence the operator

$$\Phi_{J,\delta} : Z \rightarrow W = Z - T_{CG} A_{J,\delta}(Z) \overline{Z_{\bar{\zeta}}}$$

takes the space $C^k(\overline{\mathbb{D}})$ into itself. Thus, Z is J_δ -holomorphic if and only if $\Phi_{J,\delta}(Z)$ is holomorphic (in the usual sense) on $\mathbb{D}$. For sufficiently small positive δ this is an invertible operator on a

neighborhood of zero in $C^k(\overline{\mathbb{ID}})$ which establishes a one-to-one correspondence between the sets of J_δ -holomorphic and holomorphic discs in $\mathbb{B}^n$.

These considerations allow us to replace the non-linear Riemann-Hilbert problem (RH) by Bishop's equation

$$r_\delta(\Phi_{J,\delta}^{-1}(W))(\zeta) = 0, \zeta \in b\mathbb{ID} \quad (9)$$

for an unknown holomorphic function W in the unit disc.

If W is a solution of the boundary problem (9), then $Z = \Phi_{J,\delta}^{-1}(W)$ is a Bishop disc with boundary attached to E_δ . Since the manifold E_δ is biholomorphic via non-isotropic dilations to the initial manifold E , the solutions of the equation (9) allow to describe the Bishop discs attached to E .

3.3 Analysis of the Bishop equation and the geometry of Bishop discs

We begin with the description of Bishop discs attached to the model manifold E_0 in $\mathbb{C}^n$ with the standard structure J_{st} . They are the solutions of the boundary problem (9) for $\delta = 0$.

For $r > 0$ consider the ellipse $D_r := \{w \in \mathbb{C} : P(w) < r\}$ and denote by w_r the biholomorphism $w_r = w(r, \bullet) : \mathbb{ID} \longrightarrow D_r$ satisfying $w_r(0) = 0$. Then $P \circ w_r|_{b\mathbb{ID}} \equiv r$ and we set $z_2(r, \zeta) \equiv r$. This J_{st} holomorphic disc admits the lift to a Bishop disc if we set

$$z_j(r, \zeta) = SQ_j(\zeta) + ic_n, j = 3, \dots, n$$

where $c_3, \dots, c_n$ are real constants and S denotes the Schwarz integral:

$$Sf(\zeta) = \frac{1}{2\pi i} \int_{|\tau|=1} \frac{\tau + \zeta}{\tau - \zeta} f(\tau) \frac{d\tau}{\tau}$$

We obtain a family of J_{st} -holomorphic discs $(w(r, c)(\bullet), z(r, c)(\bullet))$ smoothly depending on $(n - 1)$ real parameters $(r, c_3, \dots, c_n)$, $r > 0$. Their boundaries are disjoint and form a foliation of a neighborhood of the real $(n - 2)$ dimensional submanifold

$$\Sigma := \{w = 0, z_2 = 0, \operatorname{Re} z_3 = \dots = \operatorname{Re} z_n = 0\}$$

in E . For $r = 0$ these discs degenerate to the constant map $\zeta \mapsto (0, 0, ic_3, \dots, ic_n)$. Their images form a real $(n + 1)$ submanifold $\tilde{E}$ such that $(\tilde{E}, E)$ is a smooth manifold with boundary outside the points of Σ .

We claim that in the general case of an almost complex structure J the Bishop discs have similar properties. Indeed, Let $w = 0$, $z_2 = \delta$, $z_j = \delta$, $j = 3, \dots, n$ for $\delta > 0$ be a point on the real "normal" to E . After the non-isotropic dilation Λ_δ the image of this point coincides with $p^0 := (0, 1, 1, \dots, 1)$. There exists a unique Bishop disc Z^0 of the described above family centered at p^0 ; it corresponds to the parameters $r = 1$ and $c = 0$. The parametrizing map $F : (r, c, \zeta) \longrightarrow (w(r, c)(\zeta), z(r, c)(\zeta))$ has the maximal rank when the parameters (r, c) are in a neighborhood U of the point $(1, 0)$ and $\zeta \in \mathbb{ID}$. Applying the implicit function theorem to the operator equation (9) for $\delta > 0$ small enough we obtain a holomorphic solution $W(r, c)(\bullet)$ smoothly depending on $(n - 1)$ real parameters (r, c) .

Then $Z(r, c) = \Phi_{J, \delta}^{-1}(W)$ is a family of J -holomorphic Bishop discs with boundaries attached to E_δ . This family is a small deformation of the above Bishop disc for E_0 . Since δ is small, the parametrizing map $(r, c, \zeta) \mapsto Z(r, c)(\zeta)$ has the maximal rank. So these discs form a real smooth $(n + 1)$ manifold $\tilde{E}_\delta$ with smooth boundary. Furthermore, this manifold is foliated by the Bishop discs $Z(r, c)(\bullet)$ and a neighborhood of $Z^0(b\mathbb{D})$ in E is an open subset of the boundary of $\tilde{E}_\delta$.

We proved the following statement.

Theorem 3.3 *Let p be an elliptic point of a real n dimensional submanifold E of an almost complex manifold (M, J) . Given positive k there exists a family of J -holomorphic Bishop discs for E C^k -smoothly depending on $(n - 1)$ real parameters. Thiese discs foliate a real $(n + 1)$ -dimensional submanifold $\tilde{E}$ such that $(\tilde{E}, E)$ is a C^k smooth manifold with boundary outside a real $(n - 2)$ -dimensional submanifold Σ in E .*

As we pointed out, in the case of dimension 2 a similar statement is due to H.Hofer [5], Y.Eliashberg [3] and R.Ye [12]. It seems natural to study the questions related to the regularity of the manifold $\tilde{E}$ near the point of Σ similarly to C.Kenig - S.Webster results [6, 7] in the case of an integrable complex structure.

Aknowledgements. This paper was done while the first named author visited the University of Illinois at Urbana-Champaign during the Spring semester of 2005. He thanks this institution for excellent conditions for work.

References

- [1] E.Bedford, B.Gaveau, *Envelopes of holomorphy of certain 2-spheres in $\mathbb{C}^2$* , Amer. J. Math. **105**(1983), 975-1009.
- [2] E.Bishop, *Differentiable manifolds in complex Euclidean Space*, Duke Math. J. **32**, (1965) 1-21.
- [3] Y.Eliashberg, *Filling by holomorphic discs and its applications*, London Math. Society Lecture Notes, series 151(1991), 45-67.
- [4] M.Gromov, *Pseudo-holomorphic curves in symplectic manifolds*, Invent. Math. **82** (1985), 307-347.
- [5] H.Hofer, *Pseudoholomorphic curves in symplectizations with applications to the Weinstein conjecture in dimension three*, Invent. Math. **114**(1993), 515-563.
- [6] C.Kenig, S.Webster, *The local hull of holomorphy of a surface in the space of two complex variables*, Invent. Math. **67**(1982), 1-21.
- [7] C.Kenig, S.Webster, *On the hull of holomorphy of an n -manifold in $\mathbb{C}^n$* , Ann. Scuola Norm. Super. Pisa, **11**(1984), 261-280.
- [8] N.Kruzhilin, A.Sukhov, *Attaching pseudoholomorphic discs to real submanifolds of an almost complex space*, Bull. Sci. Math. 2005.
- [9] A.Nijenhuis, W.Woolf, *Some integration problems in almost-complex and complex manifolds*, Ann. Math. **77**(1963), 429-484.
- [10] A.Tumanov, *Extending CR functions from manifolds with boundaries*, Math. Res. Letters, **2**(1995), 629-642.
- [11] I.N.Vekua, *Generalized analytic functions*, Pergamon Press , 1962.
- [12] R.Ye, *Filling by holomorphic curves in symplectic 4-manifolds*, Trans. AMS **350**(1988), 213-250.